



(I)

wherein X_1 and X_2 independently represent O or S, R_1 , R_2 and R_3 independently represent hydrogen, lower alkyl, lower alkenyl or lower alkynyl; R_4 represents the following group:



wherein Y_1 and Y_2 independently represent hydrogen, halogen or lower alkyl, and Z represents substituted or unsubstituted aryl, or the following group:



I

α / naphthyl only

wherein m is an integer of 1 to 3 and R_6 represents hydrogen, hydroxy, lower alkyl, lower alkoxy, halogen, nitro or amino, or a substituted or unsubstituted heterocyclic group selected from furyl and pyridyl; and wherein the substituted aryl and the substituted heterocyclic group have 1 to 3 independently-selected substituents selected from the group consisting of lower alkyl, hydroxy, lower alkoxy or lower alkoxy substituted with a substituent(s) selected from the group consisting of hydroxy, lower alkoxy, halogen, amino, azido, carboxy and lower alkoxycarbonyl, halogen, nitro, amino, lower alkylamino, di(lower alkyl)amino, trifluoromethyl,

Ficent
E' cent
trifluoromethoxy, benzyloxy, phenyl, phenoxy, lower alkanoyl, lower alkanoyloxy,
aroxyloxy, aralkanoyloxy, carboxy, lower alkoxycarbonyl, lower alkylcarbamoyl,
di(lower alkyl)carbamoyl, sulfo, lower alkoxysulfonyl, lower alkylsulfamoyl and
di(lower alkyl)sulfamoyl; or a pharmaceutically acceptable salt thereof, as an active
ingredient
